



PDCD10 gene

programmed cell death 10

Normal Function

The *PDCD10* gene (also known as *CCM3*) provides instructions for making a protein that appears to play a role in the structure of blood vessels. While the exact function of the PDCD10 protein is unclear, studies suggest that it works with other proteins to help strengthen the interactions between cells and limit leakage from blood vessels. This protein is also thought to be involved in pathways that signal cells to self-destruct (undergo apoptosis) when they have completed a certain number of cell divisions or accumulated errors in their DNA.

Health Conditions Related to Genetic Changes

cerebral cavernous malformation


More than a dozen mutations in the *PDCD10* gene have been identified in families with cerebral cavernous malformations, which are collections of blood vessels in the brain that are weak and prone to leakage. These mutations include a deletion of the entire gene, deletion of small segments of DNA, and changes in single DNA building blocks (nucleotides). These mutations result in an abnormal or absent PDCD10 protein. It is unclear how mutations in the *PDCD10* gene lead to the formation of cerebral cavernous malformations.

Mutations in the *PDCD10* gene account for approximately 10 percent of familial cerebral cavernous malformation cases.

Chromosomal Location

Cytogenetic Location: 3q26.1, which is the long (q) arm of chromosome 3 at position 26.1

Molecular Location: base pairs 167,683,891 to 167,735,690 on chromosome 3 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- apoptosis-related protein 15
- CCM3
- cerebral cavernous malformation 3
- PDC10_HUMAN
- TFAR15

Additional Information & Resources

GeneReviews

- Familial Cerebral Cavernous Malformation
<https://www.ncbi.nlm.nih.gov/books/NBK1293>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28PDCD10%5BTIAB%5D%29%29+OR+%28programmed+cell+death+10%5BTIAB%5D%29%29+OR+%28TFAR15%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>

OMIM

- PROGRAMMED CELL DEATH 10
<http://omim.org/entry/609118>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_PD10.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=PD10%5Bgene%5D>
- HGNC Gene Family: STRIPAK complex
<http://www.genenames.org/cgi-bin/genefamilies/set/1371>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=8761
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/11235>
- UniProt
<http://www.uniprot.org/uniprot/Q9BUL8>

Sources for This Summary

- Bergametti F, Denier C, Labauge P, Arnoult M, Boetto S, Clanet M, Coubes P, Echenne B, Ibrahim R, Irthum B, Jacquet G, Lonjon M, Moreau JJ, Neau JP, Parker F, Tremoulet M, Tournier-Lasserre E; Société Française de Neurochirurgie. Mutations within the programmed cell death 10 gene cause cerebral cavernous malformations. *Am J Hum Genet.* 2005 Jan;76(1):42-51. Epub 2004 Nov 12.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15543491>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1196432/>
- Craig HD, Günel M, Cepeda O, Johnson EW, Ptacek L, Steinberg GK, Ogilvy CS, Berg MJ, Crawford SC, Scott RM, Steichen-Gersdorf E, Sabroe R, Kennedy CT, Mettler G, Beis MJ, Fryer A, Awad IA, Lifton RP. Multilocus linkage identifies two new loci for a mendelian form of stroke, cerebral cavernous malformation, at 7p15-13 and 3q25.2-27. *Hum Mol Genet.* 1998 Nov;7(12):1851-8.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/9811928>
- Guclu B, Ozturk AK, Pricola KL, Bilguvar K, Shin D, O'Roak BJ, Gunel M. Mutations in apoptosis-related gene, PD10, cause cerebral cavernous malformation 3. *Neurosurgery.* 2005 Nov;57(5):1008-13.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16284570>
- Liquori CL, Berg MJ, Squitieri F, Ottenbacher M, Sorlie M, Leedom TP, Cannella M, Maglione V, Ptacek L, Johnson EW, Marchuk DA. Low frequency of PD10 mutations in a panel of CCM3 probands: potential for a fourth CCM locus. *Hum Mutat.* 2006 Jan;27(1):118.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16329096>
- OMIM: PROGRAMMED CELL DEATH 10
<http://omim.org/entry/609118>
- Plummer NW, Zawistowski JS, Marchuk DA. Genetics of cerebral cavernous malformations. *Curr Neurol Neurosci Rep.* 2005 Sep;5(5):391-6. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16131422>

- Verlaan DJ, Roussel J, Laurent SB, Elger CE, Siegel AM, Rouleau GA. CCM3 mutations are uncommon in cerebral cavernous malformations. *Neurology*. 2005 Dec 27;65(12):1982-3.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16380626>
 - Voss K, Stahl S, Schleider E, Ullrich S, Nickel J, Mueller TD, Felbor U. CCM3 interacts with CCM2 indicating common pathogenesis for cerebral cavernous malformations. *Neurogenetics*. 2007 Nov; 8(4):249-56. Epub 2007 Jul 27.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17657516>
-

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/PDCD10>

Reviewed: November 2012

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services